

ON $[[n, n-4, 3]]_q$ QUANTUM MDS CODES FOR ODD PRIME POWER q

RUIHU LI⁽¹⁾⁽²⁾, ZONGBEN XU⁽¹⁾

ABSTRACT. For each odd prime power q , let $4 \leq n \leq q^2 + 1$. Hermitian self-orthogonal $[n, 2, n-1]$ codes over $GF(q^2)$ with dual distance three are constructed by using finite field theory. Hence, $[[n, n-4, 3]]_q$ quantum MDS codes for $4 \leq n \leq q^2 + 1$ are obtained.

1. INTRODUCTION

The theory of quantum error-correcting codes (QECCs, for short) was established a decade ago as the primary tool for fighting decoherence in quantum computers and quantum communication system, see [21] and [22]. The most widely studied class of quantum codes are binary quantum stabilizer codes. A thorough discussion on the principles of quantum coding theory was given in [4] and [9] for binary quantum stabilizer codes. An appealing aspect of binary quantum codes is that there exist links to classical coding theory which ease the construction of good quantum codes. Following [4] and [9], many binary quantum codes are constructed from binary and quaternary classical codes, see [6-7, 13-18, 23-25].

Almost at the same time of [4], some results of binary quantum stabilizer codes were generalized to the case of non-binary quantum stabilizer codes, and characterization of non-binary quantum stabilizer codes over $GF(q)$ in term of classical codes over $GF(q^2)$ was also given which generalizes the well-known notation of additive codes over $GF(4)$ for binary case, see [19], [2] and [11] and references therein. And, many non-binary quantum stabilizer codes are constructed from classical codes over $GF(q)$ or over $GF(q^2)$, see [2-3, 5, 8, 10-11].

One central theme in quantum error-correction is the construction of quantum codes with good parameters. Except the method of constructing quantum codes from classical self-orthogonal linear codes over $GF(q)$ and self-orthogonal additive codes over $GF(q^2)$ that given in [4], [9], [19], [2] and [11], Schlingemann and Werner [20] presented another new way to construct quantum stabilizer codes by finding certain graphs (or matrices) with specific properties. A number of researchers use these methods to construct optimal quantum codes, e.g., codes with largest

Ruihu Li is with the College of Science, Xi'an Jiaotong University, Shaanxi 710049, People's Republic of China, and College of Science, Air Force Engineering University, Xi'an, Shaanxi 710051, People's Republic of China (E-mail: liruihu2008@yahoo.com.cn).

Zongben Xu is with the Institute for Information and System Sciences, Xi'an Jiaotong University, Shaanxi 710049, People's Republic of China (E-mail: zbxu@mail.xjtu.edu.cn).

This work is supported by Natural Science Foundation of China under Grant No.60573040, the National Basic Research Program of China (973 Program) under Grant No.2007CB311002, Natural Science Basic Research Plan in Shaanxi Province of China under Program No.SJ08A02.

possible k with fixed n and d . Optimal quantum codes that saturating the quantum Singleton bound received much attention.

Lemma 1.1: [12] [19](**quantum Singleton bound**) An $[[n, k, d]]_q$ quantum stabilizer codes satisfies

$$k \leq n - 2d + 2.$$

A quantum code attains the quantum Singleton bound is called a quantum maximum distance separable code or *quantum MDS code* for short. It is known that except trivial codes (codes with $d \leq 2$), there are only two binary quantum MDS codes, $[[5, 1, 3]]_2$ and $[[6, 0, 4]]_2$, see [4]. Non-binary quantum MDS codes are much complex compared with the binary case. In the simplest nontrivial case $d = 3$, despite many efforts to construct non-binary quantum MDS codes, a systematic construction for all lengths has not been achieved yet, see [2-3, 5, 8, 10-11]. Known results on non-binary quantum MDS codes are as follows: [19] proved the existence of $[[5, 1, 3]]_p$ for odd prime p , [3] proved the existence of $[[q^2 + 1, q^2 - 3, 3]]_q$ quantum MDS codes for prime power q , [6] proved the existence of $[[n, n - 2d + 2, d]]_q$ for all $3 \leq n \leq q$ and $1 \leq d \leq \frac{n}{2} + 1$, and $[[q^2, q^2 - 2d + 2, d]]_q$ for $1 \leq d \leq q$, all these three papers used self-orthogonal codes to construct quantum MDS codes. Based on graph states method, [20] proved the existence of $[[5, 1, 3]]_p$ for prime $p \geq 3$, [5] proved the existence of $[[6, 2, 3]]_p$ and $[[7, 3, 3]]_p$ for prime $p \geq 3$. With a computer search and graph states method, [10] constructed four families of quantum MDS codes $[[6, 2, 3]]_p$, $[[7, 3, 3]]_p$, $[[8, 4, 3]]_p$, and $[[8, 2, 4]]_p$, for odd $p \geq 3$.

If $d \geq 3$, [11] proved that the maximal length n of $[[n, k, d]]_q$ quantum MDS codes satisfies $n \leq q^2 + d - 2$. In this paper, we will use Hermitian self-orthogonal codes over F_{q^2} to construct q -ary quantum MDS codes of distance three, where $q = p^r$ and $p \geq 3$ is an odd prime. Our main result of this paper is as follow:

Theorem 1.1: If $q = p^r$ and $p \geq 3$ is an odd prime, then there are $[[n, n - 4, 3]]_q$ quantum MDS codes for $4 \leq n \leq q^2 + 1$.

2. PRELIMINARIES

In order to present our main result, we make some preparation on quantum codes and finite fields.

Let $GF(q^2)^n$ be the n -dimensional row space over the finite field $GF(q^2)$. For $X = (x_1, x_2, \dots, x_n)$, $Y = (y_1, y_2, \dots, y_n) \in GF(q^2)^n$, the Hermitian inner product of X and Y is defined as follow:

$$(X, Y) = x_1 y_1^q + x_2 y_2^q + \dots + x_n y_n^q.$$

If $\mathcal{C}$ is an $[n, k]$ linear code over $GF(q^2)$, its dual code $\mathcal{C}^\perp = \{X \mid X \in GF(q^2)^n, (X, Y) = 0 \text{ for any } Y \in \mathcal{C}\}$. $\mathcal{C}$ is self-orthogonal if $\mathcal{C} \subseteq \mathcal{C}^\perp$, and self-dual if $\mathcal{C} = \mathcal{C}^\perp$.

[6] gave the following theorem of constructing quantum codes from self-orthogonal codes over $GF(q^2)$.

Theorem 2.1: If $\mathcal{C}$ is an $[n, k]$ linear code over $GF(q^2)$ such that $\mathcal{C}^\perp \subseteq \mathcal{C}$, and $d = \min\{wt(v) : v \in \mathcal{C} \setminus \mathcal{C}^\perp\}$, then there exists $[[n, 2k - n, d]]_q$ quantum code.

In the construction of self-orthogonal codes, we also need the following results on finite fields.

Lemma 2.1: If α is a primitive element of $GF(q^2)$, for each non zero element β of $GF(q)$, there are $q + 1$ elements α^i of $GF(q^2)$ such that $(\alpha^i)^{q+1} = \beta$.

Proof: Suppose $(\alpha)^{q+1} = \gamma$, then γ is a primitive element of $GF(q)$. Let $\beta = \gamma^i$, $0 \leq i \leq q - 2$. Then $(\alpha^{i+(q-1)j})^{q+1} = \beta$ for $0 \leq j \leq q$, thus the Lemma holds.

Lemma 2.2: If α is a primitive element of $GF(q^2)$, then $1 + (\alpha)^{q+1} + (\alpha^2)^{q+1} + \dots + (\alpha^{q^2-2})^{q+1} = 0$.

Proof: Suppose $(\alpha)^{q+1} = \gamma$, then $1 + (\alpha)^{q+1} + (\alpha^2)^{q+1} + \dots + (\alpha^{q^2-2})^{q+1} = (q+1)(1 + \gamma + \gamma^2 + \dots + \gamma^{q-2}) = 0$

To simplify statement of the following two section, we divide the non-zero elements of $GF(q^2)$ into two subsets, say A and B . Let $A = \{1, -1, \alpha^{\frac{q-1}{2}}, -\alpha^{\frac{q-1}{2}}, \alpha^{\frac{q-1}{2}} + 1, -\alpha^{\frac{q-1}{2}} - 1\}$, $B = \{x_1, -x_1, \dots, x_k, -x_k\} = GF(q^2) \setminus (A \cup \{0\})$, where $2k = q^2 - 7$.

According to Lemma 2.2, one can deduce that

Lemma 2.3: Suppose A and B be defined as above, then $2\sum_{i=1}^k (x_i)^{q+1} + 2(\alpha^{\frac{q-1}{2}} + 1)^{q+1} = 0$.

3. $[[n, n-4, 3]]_q$ FOR $q = 3^r$

In this section, we will prove Theorem 1.1 holds for $q = 3^r$.

First we discuss the construction of $[[n, n-4, 3]]_3$ quantum code.

Let $GF(3) = \{0, 1, 2\} = \{0, 1, -1\}$ be the Galois field with three elements. then $f(x) = x^2 + x + 2$ is irreducible over $GF(3)$. Using $f(x)$, one can construct the Galois field $GF(9)$ with nine elements as $GF(9) = \{0, 1, 2, \alpha, \alpha+1, \alpha+2, 2\alpha, 2\alpha+1, 2\alpha+2\}$, where α is a root of $f(x) = x^2 + x + 2$. It is easy to check that α is a primitive element of $GF(9)$, $\alpha^2 = 2\alpha + 1$, $\alpha^3 = 2\alpha + 2$, $\alpha^4 = 2$, $\alpha^5 = 2\alpha$, $\alpha^6 = \alpha + 2$, and $\alpha^7 = \alpha + 1$. It is obvious that $\alpha^{4+i} = -\alpha^i$ for $0 \leq i \leq 7$.

Construct

$$\begin{aligned} H_{2,4} &= \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & -1 & 1 \end{pmatrix}, H_{2,5} = \begin{pmatrix} 1 & 1 & \alpha & \alpha & 0 \\ 0 & 1 & \alpha^2 & \alpha^3 & \alpha \end{pmatrix}, \\ H_{2,6} &= \begin{pmatrix} 1 & \alpha & \alpha & \alpha & \alpha & 0 \\ 0 & 1 & \alpha^2 & \alpha^4 & \alpha^6 & \alpha \end{pmatrix}, H_{2,7} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & 1 & \alpha^1 & \alpha^2 & \alpha^5 & \alpha^7 & 1 \end{pmatrix}, \\ H_{2,8} &= \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & \alpha & \alpha & 0 \\ 0 & 1 & 2 & \alpha^1 & \alpha^5 & 1 & 2 & 1 \end{pmatrix}, H_{2,9} = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ 0 & 1 & \alpha^1 & \dots & \alpha^7 \end{pmatrix}, \\ H_{2,10} &= \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & \alpha & \alpha & \alpha & 0 \\ 0 & 1 & \alpha & \alpha^4 & \alpha^6 & \alpha^7 & \alpha^3 & \alpha^4 & \alpha^6 & 1 \end{pmatrix}, \end{aligned}$$

It is easy to check that: For $4 \leq n \leq 10$, the code $\mathcal{C}_{2,n}$ generated by $H_{2,n}$ is a self-orthogonal code over $GF(9)$ and $d^\perp = 3$, hence there is an $[[n, n-4, 3]]_3$ quantum MDS codes.

Second, we discuss the construction of $[[n, n-4, 3]]_q$ quantum code for $q = 3^r \geq 9$. To achieve this, we consider four cases separately.

Case 3.1 Let $4 \leq n \leq q^2 - 4$ and $n \equiv 0 \pmod{2}$.

Let $n - 4 = 2k_1$, $u = 2\sum_{i=1}^{k_1} (x_i)^{q+1}$. Choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\delta^{q+1} \in GF(q) \setminus GF(3)$ and $2\delta^{q+1} + u \neq 0$, $\gamma^{q+1} = -(n-4+2\delta^{q+1})$, $\epsilon^{q+1} = -(u+2\delta^{q+1})$.

Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \dots & 1 & 1 & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \dots & x_{k_1} & -x_{k_1} & \delta & -\delta & \epsilon \end{pmatrix}.$$

Case 3.2 Let $4 \leq n \leq q^2 - 4$ and $n \equiv 1 \pmod{2}$.

Let $n - 5 = 2k_1$, $u = 2\sum_{i=1}^{k_1} (x_i)^{q+1}$. Choose $\gamma, \delta, \epsilon \in GF(q^2)$, such that $\delta^{q+1} \in GF(q) \setminus GF(3)$ and $u + \delta^{q+1}(\alpha^{q-1} + 1)^{q+1} \neq 0$, $\gamma^{q+1} = -(n-5 + \delta^{q+1})$, $\epsilon^{q+1} =$

$-(u + \delta^{q+1}(\alpha^{q-1} + 1)^{q+1})$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta\alpha^{\frac{q-1}{2}} & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta\alpha^{\frac{q-1}{2}} & -\delta\alpha^{q-1} & \delta(\alpha^{q-1} + 1) & \epsilon \end{pmatrix}.$$

Case 3.3 Let $q^2 - 3 \leq n \leq q^2 - 1$.

Subcase 3.3.1 If $n = q^2 - 1$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\delta^{q+1} \in GF(q) \setminus GF(3)$ and $1 - \delta^{q+1} - (\alpha^{q-1} + 1)^{q+1} \neq 0$, $\gamma^{p+1} = 5 - 2\delta^{q+1}$, $\epsilon^{q+1} = 2(\delta^{q+1} + (\alpha^{\frac{q-1}{2}} + 1)^{q+1} - 1)$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & 1 & -1 & \delta\alpha^{\frac{q-1}{2}} & -\delta\alpha^{\frac{q-1}{2}} & \epsilon \end{pmatrix}.$$

Subcase 3.3.2 If $n = q^2 - 2$, choose $\epsilon \in GF(q^2)$ such that $\epsilon = \alpha^{\frac{q-1}{2}} + 1$, and construct

$$A_{2,n} = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & 1 & \alpha^{\frac{q-1}{2}} & -\alpha^{\frac{q-1}{2}} - 1 & \epsilon \end{pmatrix}.$$

Subcase 3.3.3 If $n = q^2 - 3$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\delta^{q+1} \in GF(q) \setminus GF(3)$, $\gamma^{p+1} = 1 - 2\delta^{q+1}$, $\epsilon^{q+1} = (2 - 2\delta^{q+1})(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & \delta(\alpha^{\frac{p-1}{2}} + 1) & -\delta(\alpha^{\frac{p-1}{2}} + 1) & \epsilon \end{pmatrix}.$$

Case 3.4 Let $q^2 \leq n \leq q^2 + 1$.

If $n = q^2$, construct

$$A_{2,n} = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 0 & 1 & \alpha & \cdots & \alpha^{q^2-3} & \alpha^{q^2-2} \end{pmatrix}.$$

If $n = q^2 + 1$, let $a = \frac{q-1}{2}$. Choose $\delta, \epsilon \in GF(q^2)$, such that $\delta^{q+1} \in GF(q) \setminus GF(3)$, $\epsilon^{q+1} = (1 - \delta^{q+1})(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$, and construct

$$A_{2,n} = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & 1 & \alpha^a & -(\alpha^a + 1) & -\delta & -\delta\alpha^a & \delta(\alpha^a + 1) & \epsilon \end{pmatrix}.$$

It is easy to check that: In the above four cases, the code generated by $A_{2,n}$ is an $[n, 2, n-1]$ self-orthogonal code over $GF(q^2)$, and its dual distance is 3. Hence, there are $[[n, n-4, 3]]_q$ quantum MDS codes for $4 \leq n \leq q^2 + 1$, where $q = 3^r$.

Summarizing the above discussion, Theorem 1.1 holds for $q = 3^r$.

4. $[[n, n-4, 3]]_q$ FOR $q = p^r$ AND PRIME $p \geq 5$

In this section, we will prove Theorem 1.1 holds for $q = p^r$, and we always assume that $p \geq 5$ is a prime and α is a primitive element of $GF(q^2)$. To give the construction of quantum $[[n, n-4, 3]]_q$ codes, we consider four cases separately.

Case 4.1 Let $4 \leq n \leq q^2 - 4$ and $n = mp + 2$.

Subcase 4.1.1 If $n \equiv 0 \pmod{2}$, let $n - 4 = 2k_1$.

If $v = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + 4 \neq 0$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -2$, $\delta^{q+1} = 2$, $\epsilon^{q+1} = -v$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta & -\delta & \epsilon \end{pmatrix}.$$

If $v = 2\Sigma_{i=1}^{k_1}(x_i)^{p+1} + 4 = 0$, choose $\gamma, \delta, \epsilon \in GF(q^2)$, such that $\gamma^{q+1} = -2$, $\delta^{q+1} = 2$, $\epsilon^{q+1} = 8$, and construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta\alpha^{\frac{q-1}{2}} & -\delta\alpha^{\frac{q-1}{2}} & \epsilon \end{pmatrix}.$$

Subcase 4.1.2 If $n \equiv 1 \pmod{2}$, let $n - 5 = 2k_1$.

If $v' = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + 2(\alpha^{\frac{q-1}{2}} + 1)^{q+1} \neq 0$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -3$, $\delta^{q+1} = 2$ and $\epsilon^{q+1} = -v'$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta & \delta\alpha^{\frac{q-1}{2}} & -\delta(\alpha^{\frac{q-1}{2}} + 1) & \epsilon \end{pmatrix}.$$

If $v' = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + 2(\alpha^{\frac{q-1}{2}} + 1)^{p+1} = 0$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -6$, $\delta^{q+1} = 3$ and $\epsilon^{q+1} = -(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta & \delta\alpha^{\frac{q-1}{2}} & -\delta(\alpha^{\frac{q-1}{2}} + 1) & \epsilon \end{pmatrix}.$$

Case 4. 2 Let $4 \leq n \leq q^2 - 4$ and $n - 2 \not\equiv 0 \pmod{p}$.

Subcase 4.2.1 If $n \equiv 0 \pmod{2}$, let $n - 4 = 2k_1$.

If $w = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + 2 \neq 0$, choose $\gamma, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -n + 2$ and $\epsilon^{q+1} = -w$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & 1 & -1 & \epsilon \end{pmatrix}.$$

If $w = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + 2 = 0$, choose $\gamma, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -n + 2$ and $\epsilon^{q+1} = 4$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \alpha^{\frac{q-1}{2}} & -\alpha^{\frac{q-1}{2}} & \epsilon \end{pmatrix}.$$

Subcase 4.2.2 If $n \equiv 1 \pmod{2}$, let $n - 5 = 2k_1$.

If $w' = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + (\alpha^{\frac{q-1}{2}} + 1)^{q+1} \neq 0$, choose $\gamma, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -n + 2$, $\epsilon^{q+1} = -w_1$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & 1 & \alpha^{\frac{q-1}{2}} & -\alpha^{\frac{q-1}{2}} - 1 & \epsilon \end{pmatrix}.$$

If $w' = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + (\alpha^{\frac{q-1}{2}} + 1)^{q+1} = 0$ and $n + 1 \not\equiv 0 \pmod{p}$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -n - 1$, $\delta^{q+1} = 2$ and $\epsilon^{q+1} = -(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta & \delta\alpha^{\frac{q-1}{2}} & -\delta\alpha^{\frac{q-1}{2}} - \delta & \epsilon \end{pmatrix}.$$

If $w' = 2\Sigma_{i=1}^{k_1}(x_i)^{q+1} + (\alpha^{\frac{q-1}{2}} + 1)^{q+1} = 0$ and $n + 1 \equiv 0 \pmod{p}$, choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -n - 4$, $\delta^{q+1} = 3$ and $\epsilon^{q+1} = -2(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$.

Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_{k_1} & -x_{k_1} & \delta & \delta\alpha^{\frac{q-1}{2}} & -\delta\alpha^{\frac{q-1}{2}} - \delta & \epsilon \end{pmatrix}.$$

Case 4.3 Let $q^2 - 3 \leq n \leq q^2 - 1$.

Subcase 4.3.1 If $n = q^2 - 1$, choose $\gamma, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = 3$, $\epsilon^{q+1} = 2(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$, and construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & 1 & -1 & \alpha^{\frac{q-1}{2}} & -\alpha^{\frac{q-1}{2}} & \epsilon \end{pmatrix}.$$

Subcase 4.3.2 If $n = q^2 - 2$, choose $\gamma, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = 4$, $\epsilon^{q+1} = (\alpha^{\frac{q-1}{2}} + 1)^{q+1}$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & 1 & \alpha^{\frac{q-1}{2}} & -\alpha^{\frac{q-1}{2}} - 1 & \epsilon \end{pmatrix}.$$

Subcase 4.3.3 Let $n = q^2 - 3$. Choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = 3$, $\delta^{q+1} = 2$, $\epsilon^{q+1} = -2(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$, and construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & \delta(\alpha^{\frac{q-1}{2}} + 1) & -\delta(\alpha^{\frac{q-1}{2}} + 1) & \epsilon \end{pmatrix}.$$

Case 4.4 Let $q^2 \leq n \leq q^2 + 1$.

If $n = q^2$, construct

$$A_{2,n} = \begin{pmatrix} 1 & 1 & 1 & 1 & \cdots & 1 & 1 \\ 0 & 1 & \alpha & \alpha^2 & \cdots & \alpha^{p^2-3} & \alpha^{p^2-2} \end{pmatrix}.$$

If $n = q^2 + 1$, let $a = \frac{q-1}{2}$. Choose $\gamma, \delta, \epsilon \in GF(q^2)$ such that $\gamma^{q+1} = -2$, $\delta^{q+1} = 2$ and $\epsilon^{q+1} = -(\alpha^{\frac{q-1}{2}} + 1)^{q+1}$. Construct

$$A_{2,n} = \begin{pmatrix} \gamma & 1 & 1 & \cdots & 1 & 1 & 1 & 1 & 1 & \delta & \delta & \delta & 0 \\ 0 & x_1 & -x_1 & \cdots & x_k & -x_k & 1 & \alpha^a & -(\alpha^a + 1) & -\delta & -\delta\alpha^a & \delta(\alpha^a + 1) & \epsilon \end{pmatrix}.$$

It is easy to check that: In the above four cases, the code generated by $A_{2,n}$ is an $[n, 2, n-1]$ self-orthogonal code over $GF(q^2)$ with dual distance is 3. Hence, there are $[[n, n-4, 3]]_q$ quantum MDS codes for $4 \leq n \leq q^2 + 1$, where $q = p^r$ and $p \geq 5$.

Summarizing the above discussion, Theorem 1.1 holds for $q = p^r$ and $p \geq 5$ is a prime.

5. CONCLUDING REMARKS

For each odd prime power q , we have constructed an $[[n, n-4, 3]]_q$ quantum MDS codes for $4 \leq n \leq q^2 + 1$. If $q = 2^r \geq 4$, are there $[[n, n-4, 3]]_q$ quantum MDS codes for $4 \leq n \leq q^2 + 1$? If such quantum MDS codes exist, how to construct them? These need further study.

REFERENCES

1. S.A. Aly, A. Klappenecker and P.K.Sarvepalli, On Quantum and Classical BCH codes, *IEEE Trans. Inform. Theory*, **53**, pp.1183-1188, 2007.
2. A. Ashikhim and E. Knill, Non-binary quantum stabilizer codes, *IEEE Trans. Inform. Theory*, **47**, pp. 3065C3072, 2001.
3. J. Bierbrauer and Y. Edel, Quantum twisted codes, *J. Comb. Designs*, **8**, pp.174-188, 2000.
4. A.R. Calderbank, E.M. Rains, P.W. Shor and N.J.A. Sloane, Quantum error-correction via codes over $GF(4)$, *IEEE Trans. Inform. Theory*, **44**, pp.1369-1387, 1998.
5. Keqin Feng, Quantum codes $[[6, 2, 3]]_p$ and $[[7, 3, 3]]_p$ ($p \geq 3$) exist, *IEEE Trans. Inform. Theory*, **48**, pp. 2384-2391, 2002.
6. M. Grassl and T. Beth, Quantum BCH codes, quant-ph/9910060, 1999.
7. M. Grassl, W. Geiselmann and T. Beth, Quantum Reed-Solomon codes, in Proc. AAECC Conf., 1999. quant-ph/9910059, 1999.
8. M. Grassl, T. Beth, and M. Rotteler. On optimal quantum codes. *Internat. J. Quantum Information*, 2(1):757-775, 2004.
9. D. Gottesman, Stabilizer codes and quantum error correction, Ph.D. Thesis, California Institute of Technology, 1997. quant-ph/9707027, 1997.
10. Dan Hu, Weidong Tang and Meisheng Zhao, et. al. Graphical nonbinary quantum error-correcting codes *Phys. Rev. A.*, **78**, 012306. 2008.
11. A. Ketkar, A. Klappenecker and S. Kumar, nonbinary stablizer codes over finite fields, *IEEE Trans. Inform. Theory*, **52**, pp.4892-4914, 2006.
12. E. Knill and R. Laflamme, A theory of quantum error-correcting codes, *Phys. Rev. A.*, **vol.55**, pp. 900C911, 1997.
13. Ruihu Li and Xueliang Li, Binary construction of quantum codes of minimum distance three and four, *IEEE Trans. Inform. Theory*, **vol.50(6)**, pp1331-1336, 2004.
14. Ruihu Li and Xueliang Li, Quantum codes constructed from binary cyclic codes, *Int. J. Quantum Inform.*, **2**, pp.265-272, 2004.
15. Ruihu Li and Xueliang Li, Binary construction of quantum codes of minimum distance five and six, *Discrete Math.* **308**, pp.1603-1611, 2008.
16. Ruihu Li, Linear quantum codes of minimum distance three, *Int. J. Quantum Inform.*, **4**, pp.265-272, 2006.
17. Xiaoyan Lin, Quantum cyclic and cinstacyclic codes, *IEEE Trans. Inf. Theory*, **50(3)**, pp.547-549, 2004.
18. Yuena Ma, Quantum codes constructed from codes chains of quaternary self-orthogonal codes. *Journal of Air Force Engineering University (Natural Science Edition, In Chinese)*, **9**, pp.83-86, 2008.
19. E.M. Rains, Nonbinary quantum codes, *IEEE Trans. Inform. Theory*, **45**, pp.1827-1832, 1999.
20. D. Schlingemann and R. F. Werner, Quantum error-correcting codes associated with graphs, *Phys. Rev. A*, vol. 65, no. 012308, 2001. quantph/ 0012111.
21. P.W. Shor, "Scheme for reducing decoherence in quantum computer memory," *Phys. Rev. A*, vol.52, pp.R2493-2496, Oct. 1995.
22. A.M. Steane, "Error correcting codes in quantum theory," *Phys. Rev. Lett*, vol.77, pp.793-797, July 1996.
23. A.M. Steane, Quantum Reed-Muller codes, *IEEE Trans. Inf. Theory*, **45**, pp.1701-1703, 1999.
24. A.M. Steane, Enlargement of Calderbank-Shor-Steane quantum codes, *IEEE Trans. Inf. Theory*, **45**, 2492-2495, 1999.
25. A. Thangaraj and S.W. McLaughlin Quantum codes from cyclic codes over $GF(4^m)$, *IEEE Trans. Inf. Theory*, **47**, pp.2492-2495, 2001.